

AMENDMENTS TO THE CLAIMS

This listing of claims replaces all prior versions and listings of claims in the application.

Listing of Claims:

1. (Currently Amended) A method of promoting the rate of BFU-E or CFU-GM hematopoietic cell multiplication, comprising administering an effective amount of a CXCR4 antagonist to BFU-E or CFU-GM hematopoietic cells, wherein the CXCR4 antagonist comprises SDF-1[P2G] (SEQ ID NO: 1) or a fragment or an amino acid analog thereof having at least 50% identity to SDF-1[P2G] (SEQ ID NO: 1) or a fragment thereof, or comprises 3-hydroxy-2-napthoic acid.
2. (Canceled)
3. (Currently Amended) A method of increasing the circulation of BFU-E or CFU-GM hematopoietic cells in a patient in need of such treatment, comprising administering to the patient an effective amount of a CXCR4 antagonist to mobilize the BFU-E or CFU-GM hematopoietic cells from a marrow locus to a peripheral blood locus, wherein the CXCR4 antagonist comprises SDF-1[P2G] (SEQ ID NO: 1) or a fragment or an amino acid analog thereof having at least 50% identity to SDF-1[P2G] (SEQ ID NO: 1) or a fragment thereof, or comprises 3-hydroxy-2-napthoic acid.
4. (Currently Amended) The method of claim 1, further comprising introducing a heterologous nucleic acid sequence encoding SDF-1[P2G] (SEQ ID NO: 1) or a fragment or an amino acid analog thereof having at least 50% identity to SDF-1[P2G] (SEQ ID NO: 1) or a fragment thereof into the BFU-E or CFU-GM hematopoietic cells for gene therapy for promoting the rate of hematopoietic cell multiplication.
5. (Withdrawn) The method of claim 1, wherein the hematopoietic cells are ex vivo.
6. (Original) The method of claim 1, wherein the hematopoietic cells are in vivo.

7. (Canceled)
8. (Currently Amended) The method of claim 1, wherein the CXCR4 antagonist amino acid analog having at least 50% identity to SDF-1[P2G] (SEQ ID NO: 1) or a fragment thereof comprises a CXCR4 antagonist peptide substitution wherein the substituent is selected from the group consisting of proline, proline-amino acid chimera, and Bicyclic Turned Dipeptide.
9. (Currently Amended) The method of claim 8, wherein the CXCR4 antagonist peptide amino acid analog having at least 50% identity to SDF-1[P2G] (SEQ ID NO: 1) or a fragment thereof is selected from the group consisting of:

~~KGVSLSYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC~~
~~IDPKLKWIQEYLEKALN~~ (SEQ ID No. 1);

KGVSPSYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC
 DPKLKWIQEYLEKALN (SEQ ID No. 2);

KGVSLPYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC
 IDPKLKWIQEYLEKALN (SEQ ID No. 3);

KGVSLSPRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC
 DPKLKWIQEYLEKALN (SEQ ID No. 4);

KGVSLSYPCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC
 DPKLKWIQEYLEKALN (SEQ ID No. 5);

KGVS~~P~~*SYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQV
CIDPKLKWIQEYLEKALN (SEQ ID No. 6);

KGVS~~L~~P*YRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQV
CIDPKLKWIQEYLEKALN (SEQ ID No. 7);

KGVSLS**P***RCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQV
CIDPKLKWIQEYLEKALN (SEQ ID No. 8);


KGVSLS**Y**P*RCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQV
CIDPKLKWIQEYLEKALN (SEQ ID No. 9);

KGVS**Btd**YRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQV
CIDPKLKWIQEYLEKALN (SEQ ID No. 10);

KGVSLS**Btd**RCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC
IDPKLKWIQEYLEKALN (SEQ ID No. 11);


KGVSLS**Btd**CPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVC
IDPKLKWIQEYLEKALN (SEQ ID No. 12);

wherein P* =



with X= Ar, Ar-OH, alkyl and more

and Btd =



X= Alkyl, Ar, Ar-OH and more


10. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

- a) KGVSLSYRCPCRFFESH
- b) KGVSLSYRC

11. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:


KGVS PSYRC PCRFFESH	(SEQ ID No. 17)
KGVSL PYRC PCRFFESH	(SEQ ID No. 18)
KGVSL SPRC PCRFFESH	(SEQ ID No. 19)
KGVSL SYPC PCRFFESH	(SEQ ID No. 20)
KGVSP* SYRC PCRFFESH	(SEQ ID No. 21)
KGVSLP* YRC PCRFFESH	(SEQ ID No. 22)
KGVSLSP* RC PCRFFESH	(SEQ ID No. 23)
KGVSLSY P *CPCRFFESH	(SEQ ID No. 24)
KGVS Btd YRCPCRFFESH	(SEQ ID No. 25)
KGVSL Btd RCPCRFFESH	(SEQ ID No. 26)
KGVSLSB td CPCRFFESH	(SEQ ID No. 27)
KGVS PSYRC	(SEQ ID No. 28)
KGVSL PYRC	(SEQ ID No. 29)
KGVSL SPRC	(SEQ ID No. 30)
KGVSL SYPC	(SEQ ID No. 31)
KGVSP* SYRC	(SEQ ID No. 32)
KGVSLP* YRC	(SEQ ID No. 33)
KGVSLSP* RC	(SEQ ID No. 34)
KGVSLSY P *C	(SEQ ID No. 35)
KGVS Btd YRC	(SEQ ID No. 36)
KGVSL Btd RC	(SEQ ID No. 37)
KGVSLSB td C	(SEQ ID No. 38)

wherein P* =



with X= Ar, Ar-OH, alkyl and more

and Btd =




X= Alkyl, Ar, Ar-OH and more

12. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:




wherein P* =




with X= Ar, Ar-OH, alkyl and more

and Btd =




X= Alkyl, Ar, Ar-OH and more

13. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:




wherein X is a natural or unnatural amino acid linker between each of the arginines at position 8 in each sequence; and,

wherein P* =



with X= Ar, Ar-OH, alkyl and more

and Btd =



X= Alkyl, Ar, Ar-OH and more

14. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSLSYRCPCRFF-G_n-LKWIQEYLEKALN (SEQ No. 63)

KGVSLSYRCPCRFFESH-G_n-LKWIQEYLEKALN (SEQ No. 64)

wherein n is 0 or an integer from 1 to 10.

15. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSLSYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN (SEQ No. 65)

KGVSLSYRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN (SEQ No. 66)

where n is 0 or an integer from 1 to 20.

16. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSPSYRCPCRFF-GGGG-LKWIQEYLEKALN; KGVSLPYRCPCRFF-GGGG-LKWIQEYLEKALN; KGVSLSPRCPCRFF-GGGG-LKWIQEYLEKALN; KGVSLSYPCPCRFF-GGGG-LKWIQEYLEKALN; KGVSPSYRCPCRFFESH-GGGG-LKWIQEYLEKALN; KGVSLPYRCPCRFFESH-GGGG-LKWIQEYLEKALN; KGVSLSPRCPCRFFESH-GGGG-LKWIQEYLEKALN; KGVSLSYPCPCRFFESH-GGGG-LKWIQEYLEKALN; KGVSPSYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN; KGVSLPYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN; KGVSLSPRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;

KGVSLSYPCPCRFF-(CH₂)_n-LKWIQEYLEKALN; KGVSPSYRCPCRFFESH-
 (CH₂)_n-LKWIQEYLEKALN; KGVSLPYRCPCRFFESH-(CH₂)_n-
 LKWIQEYLEKALN; KGVSLSPRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSYPCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN,

wherein n is 0 or an integer from 1 to 20.


17. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSP*SYRCPCRFF-GGGG-LKWIQEYLEKALN; KGVSLP*YRCPCRFF-
 GGGG-LKWIQEYLEKALN; KGVSLSP*RCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSLSYP*CPCRFF-GGGG-LKWIQEYLEKALN; KGVSP*SYRCPCRFFESH-
 GGGG-LKWIQEYLEKALN; KGVSLP*YRCPCRFFESH-GGGG-
 LKWIQEYLEKALN; KGVSLSP*RCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLSYP*CPCRFFESH-GGGG-LKWIQEYLEKALN; KGVSP*SYRCPCRFF-
 (CH₂)_n-LKWIQEYLEKALN; KGVSLP*YRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSP*RCPCRFF-(CH₂)_n-LKWIQEYLEKALN; KGVSLSYP*CPCRFF-
 (CH₂)_n-LKWIQEYLEKALN; KGVSP*SYRCPCRFFESH-(CH₂)_n-
 LKWIQEYLEKALN; KGVSLP*YRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSP*RCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSYP*CPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;

KGVSBtdYRCPCRFF-GGGG-LKWIQEYLEKALN; KGVSLBtdRCPCRFF-
 GGGG-LKWIQEYLEKALN; KGVSLSBtdCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSBtdYRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLBtdRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLSBtdCPCRFFESH-GGGG-LKWIQEYLEKALN; KGVSBtdYRCPCRFF-
 (CH₂)_n-LKWIQEYLEKALN; KGVSLBtdRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSBtdCPCRFF-(CH₂)_n-LKWIQEYLEKALN; KGVSBtdYRCPCRFFESH-
 (CH₂)_n-LKWIQEYLEKALN; KGVSLBtdRCPCRFFESH-(CH₂)_n-
 LKWIQEYLEKALN; KGVSLSBtdCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN,


wherein n is 0 or an integer from 1 to 20 and

wherein P* =



with X= Ar, Ar-OH, alkyl and more


and Btd =



X= Alkyl, Ar, Ar-OH and more

18. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSLSYRCPCRFFGGGGLKWIQEYLEKALN



KGVSLSYRCPCRFFESHGGGGLKWIQEYLEKALN



KGVSLSYRCPCRFFGGGGLKWIQEYLEKALN



KGVSLSYRCPCRFFESHGGGGLKWIQEYLEKALN



19. (Withdrawn) A CXCR4 antagonist peptide selected from the group consisting of:

KGVSLSYRCPCRFFGGGGLKWIQEYLEKALN



KGVSLSYRCPCRFFESHGGGGLKWIQEYLEKALN



KGVSLSYRCPCRFFGGGGLKWIQEYLEKALN



KGVSLSYRCPCRFFESHGGGGLKWIQEYLEKALN



20. (Withdrawn) The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSLSYRCPCRFFGGGGSKPGVIFLTKRSRQV;

KGVSLSYRCPCRFF(CH₂)_n SKPGVIFLTKRSRQV;

KGVSLSYRCPCRFFGGGGEEWVQKYVDDLELSA;

KGVSLSYRCPCRFF(CH₂)_n EEWVQKYVDDLELSA,

where n is 0 or an integer between 1 and 20.

21. (Currently Amended) A method of treating a cancer in a patient in need of such treatment comprising administering an effective amount of a CXCR4 antagonist to the patient to promote the rate of hematopoietic cell multiplication, wherein the CXCR4 antagonist comprises SDF-1[P2G] (SEQ ID NO: 1) or a fragment or an amino acid analog thereof having at least 50% identity to SDF-1[P2G] (SEQ ID NO: 1) or a fragment thereof, or comprises 3-hydroxy-2-napthoic acid, and wherein the administering comprises treatment of the cancer.

22. (Canceled)